



EUROPEAN REGION

World Confederation
for Physical Therapy

European Region of the WCPT Statement on Physiotherapy in Primary Care

ADOPTED

General Meeting 26 – 28 April 2018
Dublin, Ireland

EUROPEAN REGION OF THE WCPT STATEMENT ON PHYSIOTHERAPY IN PRIMARY CARE

European Region of the WCPT Professional Issues WG (PI WG)

This statement has been produced by the European Region of the World Confederation for Physiotherapy (ER-WCPT) to promote the role of the Physiotherapy profession within primary care, to describe the health and economic benefits to health systems and populations of having a skilled, appropriately resourced and utilised physiotherapy workforce in primary care services, and to illustrate how different models of physiotherapy service delivery are contributing to these health and cost benefits.

Physiotherapy in Primary Care

Primary health care is a comprehensive, interdisciplinary, patient centred and community-oriented approach to health care. A primary care service is the patient's first point of entry into the health care system and the continuing focal point for all their health care needs (1).

Physiotherapists' qualifications and their ability to screen (2), diagnose and provide appropriate treatment or referral (3) for musculoskeletal, neurological cardiorespiratory, pediatric and other conditions, has positioned them as important providers of quality health care within the primary care team. Physiotherapists' participation in primary care contributes to addressing the growing demand for health services and the increasing costs associated with delivering these services across Europe. Physiotherapy is an effective, comparatively low-cost high-value option to meet this rising demand (4)(5)(6). The safety and efficacy of physiotherapy in primary care is well established (7)(8) as is its effect on reducing high and over utilization of other medical services (9).

Health and Economic Benefits on Physiotherapy in Primary Care

Models of physiotherapy service delivery vary throughout European countries, ranging from prescribed treatments by a medical doctor to direct access or self-referral, where patients access physiotherapy services directly, without the need to see a doctor first. Self-referral is the norm in countries like the Netherlands, Norway, Sweden and the UK, and self-referral to physiotherapy in primary care has been fully evaluated and is recommended by the National Institute for Health and Care Excellence (NICE) in the UK. It has been shown to reduce waiting times, put patients in control, enabling them to manage their condition and live more independently, and improve health outcomes by preventing acute problems from becoming chronic and reducing long term pain and disability (10)(11), reduce time off work (12), reduce costs for the NHS in the UK and reduce rates of medication prescribing (13).

Physiotherapy services have been implemented and studied in various settings such as general practice, orthopaedic outpatient clinics, emergency departments and primary care:

In general practice musculoskeletal conditions account for up to 30% of the GP's

caseload, rising to 50% for patients over 75 (14). Yet 85% of these do not need to see a GP and can be effectively managed by a physiotherapist, freeing millions of GP appointments for other patients each year (15). Support for direct access to physiotherapy is strongly expressed by both service users and clinicians (16). Fewer patients require multiple GP visits for their musculoskeletal disorder, referral to specialists/external examinations, sick-leave recommendations or prescriptions during the following year compared to a GP-assessed group (9).

The effectiveness of physiotherapists' management of musculoskeletal conditions within orthopaedic out patient services has been well studied; the diagnoses and triage recommendations for patients with hip and knee disorders made by physiotherapists are similar to the orthopaedic surgeons while patient satisfaction is significantly higher for physiotherapist care (17). Patients' perception of quality of care in a physiotherapist-led musculoskeletal service is significantly higher than that in a standard practice group. This model of care seems to meet patients' expectations and result in a greater intention to follow advice and instructions for self-management (18). A physiotherapist with additional prescribing and injection training makes decisions analogous to those of an orthopaedic surgeon at initial consultation for orthopaedic shoulder pain, including the safe identification of patients for subacromial injection (19). This model of physiotherapist-led service delivery is safe, effective and can be delivered in primary care settings.

The effectiveness of physiotherapists' caseload management in Emergency Departments (EDs) has been well studied; Patients attending emergency departments with certain conditions can be effectively managed by physiotherapists and referred to appropriate primary care services without the need for admission to an acute hospital. This can contribute to the management of certain conditions within primary care rather than in acute hospital. When comparing similar diagnostic groups, physiotherapists are significantly more time-efficient than ED physicians while managing a great deal of their caseload independently and safely (20). Patients with musculoskeletal problems in a primary practitioner physiotherapy service in Australia had improved waiting times and length of stay relative to Australian College of Emergency Medicine (ACEM) and Department of Health targets (21). A multi-disciplinary primary care team project, including physiotherapy, for patients with long-term conditions in the UK has been shown to reduce attendances at emergency departments by 15%, reduce emergency admissions to hospital by 55%, improve patients' function by 75% and reduce their anxiety by 50%, and reduce costs of care as compared to treatment in the acute hospital setting (22). The findings of these studies show that primary care physiotherapy provides an opportunity to shape patient-centred care, improve access and offer quality care on the most appropriate level, with associated health and economic benefits.

There is evidence to support the management of certain conditions within primary care physiotherapy services rather than in acute hospitals, and some examples are described:

The Health Information and Quality Authority (HIQA) in Ireland reviewed the evidence for performing shoulder arthroscopy for a number of shoulder conditions (23), knee arthroscopy for knee osteoarthritis (24), arthroplasty for knee osteoarthritis (25) and arthroplasty for hip osteoarthritis (26) and recommended a trial of physiotherapy in

the primary care setting for at least three months before referring for a surgical opinion as the more clinical and cost-effective management.

National Institute for Health and Care Excellence (NICE) Guidelines on low back pain management (27) recommend reassurance, advice to keep active, guidance on self-management, exercise programmes and manual therapy – these are the core skills of a primary care physiotherapist which positions them as significant members of the primary care workforce dealing with low back pain. Direct access to experienced specialist physiotherapists is effective in managing spinal conditions conservatively and identifying surgical candidates appropriately (28).

There is growing evidence that targeted exercise prescription can improve the health of people in the community living with cancer and has a cost benefit (29)(30)(31) in terms of recurrence, progression and survival (35-40). Prescribed exercise can also contribute towards better survival rates by addressing post diagnosis weight gain. Weight gain has been linked to poorer survival rates in breast cancer (41).

Falls in older people are one of the main reasons older people are admitted to emergency departments. Physiotherapy to restore and maintain functional mobility, bone health, strength and balance in older people can significantly reduce their risk of falls and their risk of injury if they fall. In 2014 the Chartered Society of Physiotherapy Falls Prevention Economic Model found that if everyone over the age of 65 at risk of falling was referred to physiotherapy within primary care this could reduce the number of patients who currently end up in emergency hospital department following a fall by 225,300, saving the UK health service £331 million every year. They calculated that every £1 spent on physiotherapy produces a £1.50 return on investment (32)

Falls and COPD among older people are two of the main reasons older people are admitted to hospital and require social care in the community. They are also two areas where primary care physiotherapy can have a major impact. Physiotherapy-run pulmonary rehabilitation classes for COPD patients in the UK resulted in these patients being less likely to be admitted to hospital. Those admitted spent less time in hospital and were 26% less likely to be readmitted following discharge (33)

A paediatric physiotherapist and an orthopaedic surgeon had substantial to almost perfect agreement for diagnosis, treatment and follow-up in children referred for gait abnormalities (34), showing that physiotherapists can contribute to the effective management of such conditions in primary care.

In November 2017 the UK Government announced plans to extend fit note certification beyond GPs to a wider group of healthcare professionals, including physiotherapists, psychiatrists and senior nurses, to better identify health conditions and treatments to help workers go back into their jobs faster, and to assist people with disabilities to enter or remain at work. Fit notes are designed to help patients develop a return to work plan tailored to their individual needs. (45)

These are examples of the conditions that can be effectively managed by physiotherapists in the primary care setting. There are many other conditions and populations that physiotherapists can manage effectively within primary care, such as

neurological conditions and women's health issues, but these are less well described in the literature.

What is Physiotherapy?

Physiotherapists are autonomous health professionals who are responsible for developing, maintaining and restoring human performance throughout the lifespan using evidence-based practice. They treat or prevent conditions associated with pain, injury, disease or other impairments. Physiotherapists empower patients and their carers to manage their condition outside clinical settings and to retain their independence.

Physiotherapy education in most European countries enables physiotherapists to practice autonomously in their assessment, diagnosis, management and discharge of patients. Trained to identify serious pathology many physiotherapists are qualified to undertake medical screening of patients and to refer accordingly. In some countries trained physiotherapists are authorized to administer injections or prescribe specific drugs.

What is the ER-WCPT?

The European Region of the World Confederation for Physiotherapy (ER-WCPT) is a non-profit, non-governmental organisation that represents the physiotherapy profession at European Level. The Organisation has a membership of 39 Physiotherapy Associations, one from each of the European countries, including all the EU Member States, EEA countries and all the EU applicant countries, representing over 500.000 physiotherapists in Europe. It is one of five regions that make up the World Confederation for Physical Therapy (WCPT), a global organisation representing physiotherapists and physical therapists worldwide. The WCPT claims exclusivity to the professional names "physical therapy" and "physiotherapy" and abbreviations of same, as the sole preserve of persons who hold qualifications approved by its member organisations. The titles are used interchangeably throughout the world and refer to the same profession. In Europe "physiotherapy" is the title most commonly used, and is therefore used in this paper to encompass both titles.

ER-WCPT is a member of the European Forum for Primary Care and supports its aim of improving the health of the population by promoting strong primary care.

References

1. American Academy of Family Physicians (<http://www.aafp.org/about/policies/all/primary-care.html>)
2. Boissonnault WG, Ross MD (2012). Physical therapists referring patients to physicians: a review of case reports and series. *J Orthop Sports Phys Ther.* 2012 May;42(5):446-54. doi: 10.2519/jospt.2012.3890. Epub 2012 Jan 25 (<https://www.ncbi.nlm.nih.gov/pubmed/22282166>)
3. Ludvigsson M, Enthoven P (2012). Evaluation of physiotherapists as primary assessors of patients with musculoskeletal disorders seeking primary health care. *Physiotherapy* 2012 June Volume 98, Issue 2, Pages 131–137. ([http://www.physiotherapyjournal.com/article/S0031-9406\(11\)00425-1/abstract](http://www.physiotherapyjournal.com/article/S0031-9406(11)00425-1/abstract))
4. Marks et al (2016). Increasing capacity for the treatment of common musculoskeletal problems: a non-inferiority RCT and economic analysis of corticosteroid injection for shoulder pain comparing a physiotherapist and orthopaedic surgeon. (<http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0162679>)
5. Mitchell JM, de Lissovo G (1997). A comparison of resource use and cost in direct access versus physician referral episodes of physical therapy. *Phys Ther.* 1997 Jan;77(1):10-8. (<https://www.ncbi.nlm.nih.gov/pubmed/8996459>)
6. Ojha HA, Snyder RS, Davenport TE (2013). Direct access compared with referred physical therapy episodes of care: a systemic review. *Phys Ther.* 2014 Jan;94(1):14-30. doi: 10.2522/ptj.20130096. Epub 2013 Sep 12. (<https://www.ncbi.nlm.nih.gov/pubmed/24029295>)
7. Mintken PE et al (2015). Direct access to physical therapy services is safe in a University student health centre setting. *J Allied Health.* 2015 Fall;44(3):164-8. (<https://www.ncbi.nlm.nih.gov/pubmed/26342614>)
8. Swinkels et al (2014). An overview of 5 years of patient self-referral for physical therapy in the Netherlands. *Phys Ther.* 2014 Dec;94 (12):1785-95. doi: 10.2522/ptj.20130309. Epub 2014 Jul 31. (<https://www.ncbi.nlm.nih.gov/pubmed/25082921>)
9. Bornhoft L, Larsson ME, thorn J (2015). Physiotherapy in primary care triage – the effects on utilization of medical services at primary health care clinics by patients and sub-groups of patients with musculoskeletal disorders: a case-control study. *Physiother Theory Pract.* 2015 Jan;31(1):45-52. doi: 10.3109/09593985.2014.932035. Epub 2014 Jul 2 (<https://www.ncbi.nlm.nih.gov/pubmed/24988315>)
10. The Chartered Society of Physiotherapy (2012). Musculoskeletal physiotherapy: patient self-referral. London: National Institute for Health and Care Excellence. (<https://www.nice.org.uk/savingsandproductivityandlocalpracticeresource?ci=http%3a%2f%2fwww.evidence.nhs.uk%2fresources%2fQIPP%2f29492%2fattachme nt%3fniceorg%3dtrue>)
11. Government S (2015). Allied Health Professionals Musculoskeletal Pathway Minimum Standards: a framework for action 2015-2016. Edinburgh: Scottish Government; 2015 (<http://www.gov.scot/Publications/2015/05/4529>)
12. Department of Health. Self-referral pilots to musculoskeletal physiotherapy and the implications for improving access to other AHP services. London: Department of Health; 2008 (www.webarchive.nationalarchives.gov.uk)
13. Holdsworth LK, Webster VS, McFadyen AK. What are the costs to NHS Scotland of self-referral to physiotherapy? Results of a national trial. *Physiotherapy.*

- 2007;93 (1):3-11
14. Briggs Timothy WR. Getting it right first time: improving the quality of orthopaedic care within the National Health Service in England. London: British Orthopaedic Association: 2015:
(http://www.gettingitrightfirsttime.com/downloads/BriggsReportA4_FIN.pdf)
 15. Ludvigsson ML, Enthoven P. Evaluation of physiotherapy as primary assessors of patients with musculoskeletal disorders seeking primary health care. *Physiotherapy*. 2012;98 (2):131-7
 16. Holdsworth LK , Webster V (2004). Direct access to physiotherapy in primary care: now? -and into the future? *Physiotherapy* June 2004 Volume 90, Issue 2. Pages 64-72 ([http://www.physiotherapyjournal.com/article/S0031-9406\(16\)30491-6/abstract](http://www.physiotherapyjournal.com/article/S0031-9406(16)30491-6/abstract))
 17. Desmeules Fetal (2013). Validation of an advanced practice physiotherapy model of care in an orthopaedic outpatient clinic. *BMC Musculoskeletal Disorders* 2013 14:162
(<https://bmcmusculoskeletdisord.biomedcentral.com/articles/10.1186/1471-2474-14-162>)
 18. Samsson KS, Bernhardsson S, Larsson M (2016). Perceived quality of physiotherapist-led orthopaedic triage compared with standard practice in primary care: a randomised controlled trial. *BMC Musculoskeletal Disorders* 2016 17: 257
(<https://bmcmusculoskeletdisord.biomedcentral.com/articles/10.1186/s12891-016-1112-x>)
 19. Marks D et al (2016). Agreement between a physiotherapist and an orthopaedic surgeon regarding management and prescription of corticosteroid injection for patients with shoulder pain. *Musculoskeletal Science and Practice*. Dec 2016 Vol 26, pages 216-222. ([http://www.mskscienceandpractice.com/article/S1356-689X\(16\)30724-X/abstract](http://www.mskscienceandpractice.com/article/S1356-689X(16)30724-X/abstract))
 20. De Gruchy A, Granger C, Gorelik A (2015). Physical Therapists as Primary Practitioners in the Emergency Department: Six-Month Prospective Practice Analysis. *Phys Ther*. 2015 Sep;95 (9):1207-16. doi: 10.2522/ptj.20130552. Epub 2015 Apr 30. (<https://www.ncbi.nlm.nih.gov/pubmed/25929528>)
 21. Gill SD, Stella J (2013). Implementation and performance evaluation of an emergency department primary practitioner physiotherapy service for patients with musculoskeletal conditions. *Emerg Med Australas* 2013 Dec;25(6):558-64
(<https://www.ncbi.nlm.nih.gov/pubmed/24118803>)
 22. Kent Community Health NHS Trust. The Human Touch, transforming Community Services in Kent. Service in the spot light: Pro-Active Care: Long Term Conditions. Maidstone: Kent Community Health NHS trust; 2013
www.kentcht.nhs.uk/EasySiteWeb/GatewayLink.aspx?allid=287493
 23. Health Information and Quality Authority 2014. Health Technology Assessment of Scheduled Procedures - Shoulder Arthroscopy.
<https://www.hiqa.ie/sites/default/files/2017-01/HIQA-Shoulder-Arthroscopy%20July-2014.pdf>
 24. Health Information and Quality Authority 2014. Health Technology Assessment of Scheduled Procedures - Knee Arthroscopy.
<https://www.hiqa.ie/sites/default/files/2017-01/HIQA-Knee-Arthroscopy-July-2014.pdf>
 25. Health Information and Quality Authority 2014. Health Technology Assessment of Scheduled Procedures – Knee Arthroplasty.

- <https://www.hiqa.ie/sites/default/files/2017-01/HIQA-Knee-Arthroplasty-July-2014.pdf>
26. Health Information and Quality Authority 2014. Health Technology Assessment of Scheduled Procedures – Hip Arthroplasty.
<https://www.hiqa.ie/sites/default/files/2017-01/HIQA-Hip-Arthroplasty-July-2014.pdf>
27. NICE Guideline NG59 (2016)
(<https://www.nice.org.uk/guidance/ng59/chapter/Recommendations>)
28. A review of the surgical conversion rate and independent management of spinal extended scope practitioners in a secondary care setting
(<http://publishing.rcseng.ac.uk/doi/abs/10.1308/rcsann.2016.0054>)
29. Storic L, Under J, Kane L, Foste C, Gtigliotti M. (2013). Does Exercise Provide a Positive Therapeutic Effect for Patients with Cancer Related Fatigue? A Systematic Review. *Rehabilitation Oncology* Vol. 31, No. 4, 2013
30. Titza C, Hummler S, Thomas M, Wiskemanna J. (2016). Physical exercise in advanced cancer patients undergoing palliative treatment. *Expert Review of Quality of Life in Cancer Care*, 2016 Vol. 1, No. 6, 433–442
<http://dx.doi.org/10.1080/23809000.2016.1251292>
31. Broderick J.M., Guinan E., O'Donnell D.M., Hussey J., Tyrrell E., Normand C. (2014) Calculating the costs of an 8-week physiotherapy-led exercise intervention in deconditioned cancer survivors in the early survivorship period (the PEACH trial). *Physiotherapy* 100(2), 182-4
32. The Chartered Society of Physiotherapy. The falls prevention economic model. London: The Chartered Society of Physiotherapy; 2014.
<http://www.csp.org/documents/falls-prevention-economic-model>
33. Seymour JM, Moore L, Jolley CJ et al. Outpatient pulmonary rehabilitation following acute exacerbations of COPD. *Thorax* 2010;65(5):423-8
34. Miller S et al (2016). Agreement between a physical therapist and an orthopaedic surgeon in children referred for gait abnormalities. *Pediatric Physical therapy: Spring 2016 – volume 28 – issue 1 – p85-92*
(http://journals.lww.com/pedpt/Abstract/2016/28010/Agreement_Between_a_Physical_Therapist_and_an.21.aspx)
35. Arem H, Moore SC, Park Y, et al. Physical activity and cancer-specific mortality in the NIH-AARP Diet and Health Study cohort. *International Journal of Cancer* 2014; 135(2):423-431.
36. Chlebowski RT. Nutrition and physical activity influence on breast cancer incidence and outcome. *Breast* 2013; 22 Suppl 2:S30-37.
37. Ibrahim EM, Al-Homaidh A. Physical activity and survival after breast cancer diagnosis: meta-analysis of published studies. *Medical Oncology* 2011; 28(3):753-765.
38. Holmes MD, Chen WY, Feskanich D, Kroenke CH, Colditz GA. Physical activity and survival after breast cancer diagnosis. *JAMA* 2005; 293(20):2479-2486.
39. Holick CN, Newcomb PA, Trentham-Dietz A, et al. Physical activity and survival after diagnosis of invasive breast cancer. *Cancer Epidemiology, Biomarkers & Prevention* 2008; 17(2):379-386.
40. Ballard-Barbash R, Friedenreich CM, Courneya KS, et al. Physical activity, biomarkers, and disease outcomes in cancer survivors: a systematic review. *Journal of the National Cancer Institute* 2012; 104(11):815-840.
41. Arem H, Pfeiffer RM, Engels EA, et al. Pre- and postdiagnosis physical activity, television viewing, and mortality among patients with colorectal cancer in the

- National Institutes of Health-AARP Diet and Health Study. *Journal of Clinical Oncology* 2015; 33(2):180-188.
42. Kenfield SA, Stampfer MJ, Giovannucci E, Chan JM. Physical activity and survival after prostate cancer diagnosis in the health professionals follow-up study. *Journal of Clinical Oncology* 2011; 29(6):726-732.
43. Bonn SE, Sjölander A, Lagerros YT, et al. Physical activity and survival among men diagnosed with prostate cancer. *Cancer Epidemiology, Biomarkers & Prevention* 2015; 24(1):57-64.
44. Bradshaw PT, Ibrahim JG, Stevens J, et al. Postdiagnosis change in bodyweight and survival after breast cancer diagnosis. *Epidemiology* 2012; 23(2):320-327.
45. <https://www.gov.uk/government/news/government-sets-out-plan-to-see-more-disabled-people-in-work>